

# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

**NAME** : Mr. BALWANT SINGH

AGE/ GENDER : 59 YRS/MALE **PATIENT ID** : 1755335

**COLLECTED BY** REG. NO./LAB NO. : 122502130011

REFERRED BY **REGISTRATION DATE** : 13/Feb/2025 10:44 AM BARCODE NO. : 12506993 **COLLECTION DATE** : 13/Feb/2025 11:34AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 13/Feb/2025 02:05PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

### CLINICAL CHEMISTRY/BIOCHEMISTRY

**CHOLESTEROL: SERUM** 

CHOLESTEROL TOTAL: SERUM OPTIMAL: < 200.0  $270.35^{H}$ mg/dL by CHOLESTEROL OXIDASE PAP

BORDERLINE HIGH: 200.0 -

239.0

HIGH CHOLESTEROL: > OR =

240.0

#### **INTERPRETATION:**

NATIONAL LIPID ASSOCIATION RECOMMENDATIONS (NLA-2014)	CHOLESTEROL IN ADULTS (mg/dL)	CHOLESTEROL IN ADULTS (mg/dL)
DESIRABLE	< 200.0	< 170.0
BORDERLINE HIGH	200.0 <del>- 239.</del> 0	171.0 – 199.0
HIGH	>= 240.0	>= 200.0

1. Measurements in the same patient can show physiological & analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per National Lipid association - 2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

# A PIONEER DIAGNOSTIC CENTRE

**NAME** : Mr. BALWANT SINGH

AGE/ GENDER : 59 YRS/MALE **PATIENT ID** : 1755335

**COLLECTED BY** : 122502130011 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 13/Feb/2025 10:44 AM BARCODE NO. : 12506993 **COLLECTION DATE** : 13/Feb/2025 11:34AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 13/Feb/2025 02:05PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

#### URIC ACID

URIC ACID: SERUM 3.66 mg/dL 3.60 - 7.70

by URICASE - OXIDASE PEROXIDASE

### **INTERPRETATION:-**

1.GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint.

2.Uric Acid is the end product of purine metabolism. Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation.

#### INCREASED:-

#### (A).DUE TO INCREASED PRODUCTION:-

1.Idiopathic primary gout.

2. Excessive dietary purines (organ meats, legumes, anchovies, etc).

3. Cytolytic treatment of malignancies especially leukemais & lymphomas.

4. Polycythemai vera & myeloid metaplasia.

5. Psoriasis.

Sickle cell anaemia etc.

#### (B).DUE TO DECREASED EXCREATION (BY KIDNEYS)

- 1. Alcohol ingestion.
- 2. Thiazide diuretics.
- 3.Lactic acidosis.
- 4. Aspirin ingestion (less than 2 grams per day ).
- 5. Diabetic ketoacidosis or starvation.
- 6.Renal failure due to any cause etc.

#### DECREASED:-

#### (A).DUE TO DIETARY DEFICIENCY

- 1. Dietary deficiency of Zinc, Iron and molybdenum.
- 2. Fanconi syndrome & Wilsons disease.
- 3. Multiple sclerosis
- 4. Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc.

#### (B).DUE TO INCREASED EXCREATION

1.Drugs:-Probenecid, sulphinpyrazone, aspirin doses (more than 4 grams per day), corticosterroids and ACTH, anti-coagulants and estrogens etc.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

# A PIONEER DIAGNOSTIC CENTRE

**NAME** : Mr. BALWANT SINGH

AGE/ GENDER : 59 YRS/MALE **PATIENT ID** : 1755335

**COLLECTED BY** : 122502130011 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 13/Feb/2025 10:44 AM BARCODE NO. : 12506993 **COLLECTION DATE** : 13/Feb/2025 11:34AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 13/Feb/2025 02:05PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

#### **CALCIUM**

CALCIUM: SERUM 9.15 mg/dL 8.50 - 10.60

by ARSENAZO III, SPECTROPHOTOMETRY

### **INTERPRETATION:-**

- 1. Serum calcium (total) estimation is used for the diagnosis and monitoring of a wide range of disorders including diseases of bone, kidney, parathyroid gland, or gastrointestinal tract.
- 2. Calcium levels may also reflect abnormal vitamin D or protein levels.
- 3. The calcium content of an adult is somewhat over 1 kg (about 2% of the body weight). Of this, 99% is present as calcium hydroxyapatite in bones and <1% is present in the extra-osseous intracellular space or extracellular space (ECS).
- 4. In serum, calcium is bound to a considerable extent to proteins (approximately 40%), 10% is in the form of inorganic complexes, and 50% is present as free or ionized calcium.

NOTE:-Calcium ions affect the contractility of the heart and the skeletal musculature, and are essential for the function of the nervous system. In addition, calcium ions play an important role in blood clotting and bone mineralization.

#### HYPOCALCEMIA (LOW CALCIUM LEVELS) CAUSES :-

- 1.Due to the absence or impaired function of the parathyroid glands or impaired vitamin-D synthesis.
- 2. Chronic renal failure is also frequently associated with hypocalcemia due to decreased vitamin-D synthesis as well as hyperphosphatemia and skeletal resistance to the action of parathyroid hormone (PTH).
- 3. NOTE:- A characteristic symptom of hypocalcemia is latent or manifest tetany and osteomalacia.

#### HYPERCALCEMIA (INCREASE CALCIUM LEVELS) CAUSES:-

- 1.Increased mobilization of calcium from the skeletal system or increased intestinal absorption.
- 2. Primary hyperparathyroidism (pHPT)
- 3. Bone metastasis of carcinoma of the breast, prostate, thyroid gland, or lung

**NOTE:**-Severe hypercalcemia may result in cardiac arrhythmia.

\*\*\* End Of Report \*\*\*



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

