



A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. RAJWANT KAUR

AGE/ GENDER : 38 YRS/FEMALE **PATIENT ID** : 1594519

COLLECTED BY REG. NO./LAB NO. : 122502210006

REFERRED BY **REGISTRATION DATE** : 21/Feb/2025 10:04 AM BARCODE NO. : 12507144 **COLLECTION DATE** : 21/Feb/2025 11:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 21/Feb/2025 01:02PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

HAEMATOLOGY HAEMOGLOBIN (HB)

12.1 HAEMOGLOBIN (HB) 12.0 - 16.0 gm/dL

by CALORIMETRIC

INTERPRETATION:-Hemoglobin is the protein molecule in red blood cells that carries oxygen from the lungs to the bodys tissues and returns carbon dioxide from the tissues back to the lungs.

A low hemoglobin level is referred to as ANEMIA or low red blood count.

ANEMIA (DECRESED HAEMOGLOBIN):

1) Loss of blood (traumatic injury, surgery, bleeding, colon cancer or stomach ulcer)

2) Nutritional deficiency (iron, vitamin B12, folate)

3) Bone marrow problems (replacement of bone marrow by cancer)

4) Suppression by red blood cell synthesis by chemotherapy drugs

5) Kidney failure

6) Abnormal hemoglobin structure (sickle cell anemia or thalassemia).

POLYCYTHEMIA (INCREASED HAEMOGLOBIN):

- 1) People in higher altitudes (Physiological)
- 2) Smoking (Secondary Polycythemia)
- 3) Dehydration produces a falsely rise in hemoglobin due to increased haemoconcentration
- 4) Advanced lung disease (for example, emphysema)
- 5) Certain tumors
- 6) A disorder of the bone marrow known as polycythemia rubra vera,

7) Abuse of the drug erythropoetin (Epogen) by athletes for blood doping purposes (increasing the amount of oxygen available to the body by chemically raising the production of red blood cells).

NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)



A PIONEER DIAGNOSTIC CENTRE

NAME : Mrs. RAJWANT KAUR

AGE/ GENDER : 38 YRS/FEMALE **PATIENT ID** : 1594519

COLLECTED BY REG. NO./LAB NO. : 122502210006

REFERRED BY **REGISTRATION DATE** : 21/Feb/2025 10:04 AM BARCODE NO. **COLLECTION DATE** : 21/Feb/2025 11:53AM : 12507144

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Value Unit **Test Name Biological Reference interval**

ERYTHROCYTE SEDIMENTATION RATE (ESR)

ERYTHROCYTE SEDIMENTATION RATE (ESR)

20 mm/1st hr

REPORTING DATE

0 - 20

: 21/Feb/2025 03:16PM

by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY

INTERPRETATION:

CLIENT CODE.

- 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
- 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
- 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus

CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

NOTE:

- 1. ESR and C reactive protein (C-RP) are both markers of inflammation.
- 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
 3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
 4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibringen.
 5. Women tend to average mathyldone and entraceptives professional processing mathyldone and with the opposition of the oppositio

- 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

NAME : Mrs. RAJWANT KAUR

AGE/ GENDER : 38 YRS/FEMALE **PATIENT ID** : 1594519

COLLECTED BY REG. NO./LAB NO. : 122502210006

REFERRED BY **REGISTRATION DATE** : 21/Feb/2025 10:04 AM BARCODE NO. : 12507144 **COLLECTION DATE** : 21/Feb/2025 11:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 21/Feb/2025 01:12PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

CLINICAL CHEMISTRY/BIOCHEMISTRY **GLUCOSE RANDOM (R)**

90.11 GLUCOSE RANDOM (R): PLASMA NORMAL: < 140.00 mg/dL

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 140.0 - 200.0 DIABETIC: > 0R = 200.0

INTERPRETATION

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A random plasma glucose level below 140 mg/dl is considered normal.

2. A random glucose level between 140 - 200 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prnadial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A random glucose level of above 200 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







A PIONEER DIAGNOSTIC CENTRE

NAME : Mrs. RAJWANT KAUR

AGE/ GENDER : 38 YRS/FEMALE **PATIENT ID** : 1594519

COLLECTED BY : 122502210006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 21/Feb/2025 10:04 AM BARCODE NO. : 12507144 **COLLECTION DATE** : 21/Feb/2025 11:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 21/Feb/2025 01:02PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

ENDOCRINOLOGY THYROID STIMULATING HORMONE (TSH)

THYROID STIMULATING HORMONE (TSH): SERUM 3.06

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

μIU/mL 0.35 - 5.50

3rd GENERATION, ULTRASENSITIVE

INTERPRETATION:

| AGE | REFFERENCE RANGE (μIU/mL) |
|---------------------|---------------------------|
| 0 – 5 DAYS | 0.70 - 15.20 |
| 6 Days – 2 Months | 0.70 - 11.00 |
| 3 – 11 Months | 0.70 - 8.40 |
| 1 – 5 Years | 0.70 - 7.00 |
| 6 – 10 Years | 0.60 - 5.50 |
| 11 - 15 | 0.50 - 5.50 |
| > 20 Years (Adults) | 0.27 - 5.50 |
| PRI | EGNANCY |
| 1st Trimester | 0.10 - 3.00 |
| 2nd Trimester | 0.20 - 3.00 |
| 3rd Trimester | 0.30 - 4.10 |

NOTE:-TSH levels are subjected to circardian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50 %. Hence time of the day has influence on the measured serum TSH concentration.

USE: TSH controls biosynthesis and release of thyroid harmones T4 & T3. It is a sensitive measure of thyroid function, especially useful in early or subclinical hypothyroidism, before the patient develops any clinical findings or goitre or any other thyroid function abnormality.

INCREASED LEVELS:

- 1. Primary or untreated hypothyroidism, may vary from 3 times to more than 100 times normal depending on degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis.
- 4.DRUGS: Amphetamines, Iodine containing agents and dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge.

DECREASED LEVELS:

- 1. Toxic multi-nodular goitre & Thyroiditis.
- 2. Over replacement of thyroid harmone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituatary or hypothalmic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

REPORTING DATE

: 21/Feb/2025 01:02PM

NAME : Mrs. RAJWANT KAUR

AGE/ GENDER : 38 YRS/FEMALE **PATIENT ID** : 1594519

COLLECTED BY REG. NO./LAB NO. : 122502210006

REFERRED BY **REGISTRATION DATE** : 21/Feb/2025 10:04 AM BARCODE NO. : 12507144 **COLLECTION DATE** : 21/Feb/2025 11:53AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name Value Unit **Biological Reference interval**

8. Pregnancy: 1st and 2nd Trimester LIMITATIONS:

CLIENT CODE.

1.TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

2. Autoimmune disorders may produce spurious results.

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. RAJWANT KAUR

AGE/ GENDER : 38 YRS/FEMALE **PATIENT ID** : 1594519

COLLECTED BY REG. NO./LAB NO. : 122502210006

REFERRED BY **REGISTRATION DATE** : 21/Feb/2025 10:04 AM BARCODE NO. : 12507144 **COLLECTION DATE** : 21/Feb/2025 11:53AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 21/Feb/2025 04:11PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval Test Name**

IMMUNOPATHOLOGY/SEROLOGY **C-REACTIVE PROTEIN (CRP)**

C-REACTIVE PROTEIN (CRP) QUANTITATIVE: 0.32 0.0 - 6.0

SERUM by NEPHLOMETRY

INTERPRETATION:

C-reactive protein (CRP) is one of the most sensitive acute-phase reactants for inflammation.

2. CRP levels can increase dramatically (100-fold or more) after severe trauma, bacterial infection, inflammation, surgery, or neoplastic

3. CRP levels (Quantitative) has been used to assess activity of inflammatory disease, to detect infections after surgery, to detect transplant

rejection, and to monitor these inflammatory processes.

4. As compared to ESR, CRP shows an earlier rise in inflammatory disorders which begins in 4-6 hrs, the intensity of the rise being higher than ESR and the recovery being earlier than ESR. Unlike ESR, CRP levels are not influenced by hematologic conditions like Anemia, Polycythemia etc., 5. Elevated values are consistent with an acute inflammatory process.

NOTE:

1. Elevated C-reactive protein (CRP) values are nonspecific and should not be interpreted without a complete clinical history.

2. Oral contraceptives may increase CRP levels.

*** End Of Report ***



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

