



P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

TEST PERFORMED AT KOS DIAGNOSTIC LAB, AMBALA CANTT.

NAME	: Mrs. MAYA DEVI	PATIENT ID	: 1766000
AGE/ GENDER	: 51 YRS/FEMALE	REG. NO./LAB NO.	: 122502220003
COLLECTED BY	:	REGISTRATION DATE	: 22/Feb/2025 08:44 AM
REFERRED BY	:	COLLECTION DATE	: 22/Feb/2025 09:09AM
BARCODE NO.	: 12507160	REPORTING DATE	: 22/Feb/2025 11:36AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

HAEMATOLOGY

ERYTHROCYTE SEDIMENTATION RATE (ESR)

ERYTHROCYTE SEDIMENTATION RATE (ESR)	19	mm/1st hr	0 - 20
--------------------------------------	----	-----------	--------

by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY

INTERPRETATION:

1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus

CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

NOTE:

1. ESR and C - reactive protein (C-RP) are both markers of inflammation.
2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
3. **CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.**
4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibrinogen.
5. Women tend to have a higher ESR, and menstruation and pregnancy can cause temporary elevations.
6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME	: Mrs. MAYA DEVI	PATIENT ID	: 1766000
AGE/ GENDER	: 51 YRS/FEMALE	REG. NO./LAB NO.	: 122502220003
COLLECTED BY	:	REGISTRATION DATE	: 22/Feb/2025 08:44 AM
REFERRED BY	:	COLLECTION DATE	: 22/Feb/2025 09:09AM
BARCODE NO.	: 12507160	REPORTING DATE	: 22/Feb/2025 11:36AM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE		
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

CLINICAL CHEMISTRY/BIOCHEMISTRY

URIC ACID

URIC ACID: SERUM by URICASE - OXIDASE PEROXIDASE	6.34	mg/dL	2.50 - 6.80
---	------	-------	-------------

INTERPRETATION:-

1. GOUT occurs when high levels of Uric Acid in the blood cause crystals to form & accumulate around a joint.
2. Uric Acid is the end product of purine metabolism. Uric acid is excreted to a large degree by the kidneys and to a smaller degree in the intestinal tract by microbial degradation.

INCREASED:-

(A).DUE TO INCREASED PRODUCTION:-

1. Idiopathic primary gout.
2. Excessive dietary purines (organ meats, legumes, anchovies, etc).
3. Cytolytic treatment of malignancies especially leukemias & lymphomas.
4. Polycythemia vera & myeloid metaplasia.
5. Psoriasis.
6. Sickle cell anaemia etc.

(B).DUE TO DECREASED EXCRETION (BY KIDNEYS)

1. Alcohol ingestion.
2. Thiazide diuretics.
3. Lactic acidosis.
4. Aspirin ingestion (less than 2 grams per day).
5. Diabetic ketoacidosis or starvation.
6. Renal failure due to any cause etc.

DECREASED:-

(A).DUE TO DIETARY DEFICIENCY

1. Dietary deficiency of Zinc, Iron and molybdenum.
2. Fanconi syndrome & Wilson's disease.
3. Multiple sclerosis.
4. Syndrome of inappropriate antidiuretic hormone (SIADH) secretion & low purine diet etc.

(B).DUE TO INCREASED EXCRETION

1. Drugs:- Probenecid, sulfinpyrazone, aspirin doses (more than 4 grams per day), corticosteroids and ACTH, anti-coagulants and estrogens etc.



DR. VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR. YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrajainhealthcare@gmail.com

NAME : Mrs. MAYA DEVI
AGE/ GENDER : 51 YRS/FEMALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 12507160
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

PATIENT ID : 1766000
REG. NO./LAB NO. : 122502220003
REGISTRATION DATE : 22/Feb/2025 08:44 AM
COLLECTION DATE : 22/Feb/2025 09:09AM
REPORTING DATE : 22/Feb/2025 01:11PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

IMMUNOPATHOLOGY/SEROLOGY

C-REACTIVE PROTEIN (CRP)

C-REACTIVE PROTEIN (CRP) QUANTITATIVE: 2.92 mg/L 0.0 - 6.0

SERUM

by NEPHLOMETRY

INTERPRETATION:

1. C-reactive protein (CRP) is one of the most sensitive acute-phase reactants for inflammation.
2. CRP levels can increase dramatically (100-fold or more) after severe trauma, bacterial infection, inflammation, surgery, or neoplastic proliferation.
3. CRP levels (Quantitative) has been used to assess activity of inflammatory disease, to detect infections after surgery, to detect transplant rejection, and to monitor these inflammatory processes.
4. As compared to ESR, CRP shows an earlier rise in inflammatory disorders which begins in 4-6 hrs, the intensity of the rise being higher than ESR and the recovery being earlier than ESR. Unlike ESR, CRP levels are not influenced by hematologic conditions like Anemia, Polycythemia etc.,
5. Elevated values are consistent with an acute inflammatory process.

NOTE:

1. Elevated C-reactive protein (CRP) values are nonspecific and should not be interpreted without a complete clinical history.
2. Oral contraceptives may increase CRP levels.



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)





P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

☎ 0171-2532620, 8222896961 ✉ pkrjainhealthcare@gmail.com

NAME : Mrs. MAYA DEVI
AGE/ GENDER : 51 YRS/FEMALE
COLLECTED BY :
REFERRED BY :
BARCODE NO. : 12507160
CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE
CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

PATIENT ID : 1766000
REG. NO./LAB NO. : 122502220003
REGISTRATION DATE : 22/Feb/2025 08:44 AM
COLLECTION DATE : 22/Feb/2025 09:09AM
REPORTING DATE : 22/Feb/2025 01:11PM

Test Name	Value	Unit	Biological Reference interval
-----------	-------	------	-------------------------------

RHEUMATOID FACTOR (RA): QUANTITATIVE - SERUM

RHEUMATOID (RA) FACTOR QUANTITATIVE: SERUM by NEPHLOMETRY	8.84	IU/mL	NEGATIVE: < 18.0 BORDERLINE: 18.0 - 25.0 POSITIVE: > 25.0
---	------	-------	---

INTERPRETATION:-

RHEUMATOID FACTOR (RA):

1. Rheumatoid factors (RF) are antibodies that are directed against the Fc fragment of IgG altered in its tertiary structure.
2. Over 75% of patients with rheumatoid arthritis (RA) have an IgM antibody to IgG immunoglobulin. This autoantibody (RF) is diagnostically useful although it may not be etiologically related to RA.
3. Inflammatory Markers such as ESR & C-Reactive protein (CRP) are normal in about 60 % of patients with positive RA.
4. The titer of RF correlates poorly with disease activity, but those patients with high titers tend to have more severe disease course.
5. The test is useful for diagnosis and prognosis of rheumatoid arthritis.

RHEUMATOID ARTHRITIS:

1. Rheumatoid Arthritis is a systemic autoimmune disease that is multi-functional in origin and is characterized by chronic inflammation of the membrane lining (synovium) joints which leads to progressive joint destruction and in most cases to disability and reduction of quality life.
2. The disease spreads from small to large joints, with greatest damage in early phase.
3. The diagnosis of RA is primarily based on clinical, radiological & immunological features. The most frequent serological test is the measurement of RA factor.

CAUTION (FALSE POSTIVE):-

1. RA factor is not specific for Rheumatoid arthritis, as it is often present in healthy individuals with other autoimmune diseases and chronic infections.
2. Non rheumatoid and rheumatoid arthritis (RA) populations are not clearly separate with regard to the presence of rheumatoid factor (RF) (15% of RA patients have a nonreactive titer and 8% of nonrheumatoid patients have a positive titer).
3. Patients with various nonrheumatoid diseases characterized by chronic inflammation may have positive tests for RF. These diseases include systemic lupus erythematosus, polymyositis, tuberculosis, syphilis, viral hepatitis, infectious mononucleosis, and influenza.
4. Anti-CCP have been discovered in joints of patients with RA, but not in other form of joint disease. Anti-CCP2 is HIGHLY SENSITIVE (71%) & more specific (98%) than RA factor.
5. Upto 30 % of patients with Seronegative Rheumatoid arthritis also show Anti-CCP antibodies.
6. The positive predictive value of Anti-CCP antibodies for Rheumatoid Arthritis is far greater than Rheumatoid factor.

*** End Of Report ***



DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY)

