P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

0171-2532620, 8222896961 pkrjainhealthcare@gmail.com

NAME	**: Mr. JITENDER SINGH**		
AGE/ GENDER	: 44 YRS/MALE	**PATIENT ID**	: 1539128
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122503170018**
REFERRED BY	:	**REGISTRATION DATE**	: 17/Mar/2025 11:19 AM
BARCODE NO.	: 12507545	**COLLECTION DATE**	: 17/Mar/2025 12:19PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 17/Mar/2025 02:27PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

CLINICAL CHEMISTRY/BIOCHEMISTRY

GLUCOSE FASTING (F)

GLUCOSE FASTING (F): PLASMA *by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD)*	**110.04^{H}**	mg/dL	NORMAL: < 100.0 PREDIABETIC: 100.0 - 125.0 DIABETIC: > 0R = 126.0

INTERPRETATION

IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.
2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood test (after consumption of 75 gms of glucose) is recommended for all such patients.
3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mr. JITENDER SINGH**		
AGE/ GENDER	: 44 YRS/MALE	**PATIENT ID**	: 1539128
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122503170018**
REFERRED BY	:	**REGISTRATION DATE**	: 17/Mar/2025 11:19 AM
BARCODE NO.	: 12507545	**COLLECTION DATE**	: 17/Mar/2025 12:19PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 17/Mar/2025 02:27PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

LIPID PROFILE : BASIC

Test Name	Value	Unit	Biological Reference interval
CHOLESTEROL TOTAL: SERUM *by CHOLESTEROL OXIDASE PAP*	159.82	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: SERUM *by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC)*	67.09	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTEROL (DIRECT): SERUM *by SELECTIVE INHIBITION*	41.84	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROL: SERUM *by CALCULATED, SPECTROPHOTOMETRY*	104.56	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLESTEROL: SERUM *by CALCULATED, SPECTROPHOTOMETRY*	117.98	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTEROL: SERUM *by CALCULATED, SPECTROPHOTOMETRY*	13.42	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SERUM *by CALCULATED, SPECTROPHOTOMETRY*	386.73	mg/dL	350.00 - 700.00
CHOLESTEROL/HDL RATIO: SERUM *by CALCULATED, SPECTROPHOTOMETRY*	3.82	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0 HIGH RISK: > 11.0

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

P K R JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

0171-2532620, 8222896961 pkrjainhealthcare@gmail.com

NAME	**: Mr. JITENDER SINGH**		
AGE/ GENDER	: 44 YRS/MALE	**PATIENT ID**	: 1539128
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122503170018**
REFERRED BY	:	**REGISTRATION DATE**	: 17/Mar/2025 11:19 AM
BARCODE NO.	: 12507545	**COLLECTION DATE**	: 17/Mar/2025 12:19PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 17/Mar/2025 02:27PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
LDL/HDL RATIO: SERUM *by CALCULATED, SPECTROPHOTOMETRY*	2.5	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0
TRIGLYCERIDES/HDL RATIO: SERUM *by CALCULATED, SPECTROPHOTOMETRY*	**1.6^{L}**	RATIO	3.00 - 5.00

INTERPRETATION:

1.Measurements in the same patient can show physiological& analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL , VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non HDL.

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mr. JITENDER SINGH**		
AGE/ GENDER	: 44 YRS/MALE	**PATIENT ID**	: 1539128
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122503170018**
REFERRED BY	:	**REGISTRATION DATE**	: 17/Mar/2025 11:19 AM
BARCODE NO.	: 12507545	**COLLECTION DATE**	: 17/Mar/2025 12:19PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 17/Mar/2025 04:47PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval
KIDNEY FUNCTION TEST (BASIC)			
UREA: SERUM *by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)*	26.93	mg/dL	10.00 - 50.00
CREATININE: SERUM *by ENZYMATIC, SPECTROPHOTOMETERY*	1.08	mg/dL	0.40 - 1.40
BLOOD UREA NITROGEN (BUN): SERUM *by CALCULATED, SPECTROPHOTOMETERY*	12.58	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM *by CALCULATED, SPECTROPHOTOMETERY*	11.65	RATIO	10.0 - 20.0
UREA/CREATININE RATIO: SERUM *by CALCULATED, SPECTROPHOTOMETERY*	24.94	RATIO	
URIC ACID: SERUM *by URICASE - OXIDASE PEROXIDASE*	3.62	mg/dL	3.60 - 7.70

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)

NAME	**: Mr. JITENDER SINGH**		
AGE/ GENDER	: 44 YRS/MALE	**PATIENT ID**	: 1539128
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122503170018**
REFERRED BY	:	**REGISTRATION DATE**	: 17/Mar/2025 11:19 AM
BARCODE NO.	: 12507545	**COLLECTION DATE**	: 17/Mar/2025 12:19PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 17/Mar/2025 04:47PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

INTERPRETATION:

Normal range for a healthy person on normal diet: 12 - 20

To Differentiate between pre- and postrenal azotemia.

INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1.Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion,dehydration, blood loss) due to decreased glomerular filtration rate.
2.Catabolic states with increased tissue breakdown.
3.GI hemorrhage.
4.High protein intake.
5.Impaired renal function plus .
6.Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushings syndrome, high protein diet, burns,surgery, cachexia, high fever).
7.Urine reabsorption (e.g. ureterocolostomy)
8.Reduced muscle mass (subnormal creatinine production)
9.Certain drugs (e.g. tetracycline, glucocorticoids)

INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

1.Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
2.Prerenal azotemia superimposed on renal disease.

DECREASED RATIO (<10:1) WITH DECREASED BUN :

1.Acute tubular necrosis.
2.Low protein diet and starvation.
3.Severe liver disease.
4.Other causes of decreased urea synthesis.
5.Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
6.Inherited hyperammonemias (urea is virtually absent in blood).
7.SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.
8.Pregnancy.

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:

1.Phenacimide therapy (accelerates conversion of creatine to creatinine).
2.Rhabdomyolysis (releases muscle creatinine).
3.Muscular patients who develop renal failure.

INAPPROPIATE RATIO:

1.Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies,resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).
2.Cephalosporin therapy (interferes with creatinine measurement).

*** End Of Report ***

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)