

CULTUDE AND CUCCEDTIDIL PTV. UDINE

PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

0171-2532620, 8222896961 🛛 🖾 pkrjainhealthcare@gmail.com

Test Name	Val	ue Unit	Biological Reference interval
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CI	IY - HARYANA	
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	REPORTING DATE	: 21/Mar/2025 03:17PM
BARCODE NO.	: 12507583	COLLECTION DATE	: 19/Mar/2025 10:16AM
REFERRED BY	:	REGISTRATION DATE	: 19/Mar/2025 09:45 AM
COLLECTED BY	:	REG. NO./LAB NO.	: 122503190007
AGE/ GENDER	: 22 YRS/MALE	PATIENT ID	: 1679001
NAME	: Mr. ARUN		

CULTURE AEROBIC BACTERIA AND ANTIBIOTIC SENSITIVITY: URINE

CULTURE AND SUSCEPTIBILITY: URIN	
DATE OF SAMPLE	19-03-2025
SPECIMEN SOURCE	URINE
INCUBATION PERIOD by AUTOMATED BROTH CULTURE	48 HOURS
CULTURE by AUTOMATED BROTH CULTURE	STERILE
ORGANISM by AUTOMATED BROTH CULTURE	N <mark>O AEROBIC</mark> PYOGENIC ORGANISM GROWN AFTER 48 HOURS OF INCUBATION AT 37*C
AEROBIC SUSCEPTIBILITY: URINE	

INTERPRETATION:

1. In urine culture and sensitivity, presence of more than 100,000 organism per mL in midstream sample of urine is considered clinically significant. However in symptomatic patients, a smaller number of bacteria (100 to 10000/mL) may signify infection. 2. Colony count of 100 to 10000/ mL indicate infection, if isolate from specimen obtained by suprapubic aspiration or "in-and-out" catheterization or from patients with indwelling catheters.

SUSCEPTIBILITY:

1. A test interpreted as SENSTITIVE implies that infection due to isolate may be appropriately treated with the dosage of an antimicrobial agent recommended for that type of infection and infecting species, unless otherwise indicated.. 2. A test interpreted as **INTERMEDIATE** implies that the" Infection due to the isolate may be appropriately treated in body sites where the drugs are

physiologically concentrated or when a high dosage of drug can be used". 3.A test interpreted as **RESISTANT** implies that the "isolates are not inhibited by the usually achievable concentration of the agents with normal

dosage, schedule and/or fall in the range where specific microbial resistance mechanism are likely (e.g. beta-lactamases), and clinical efficacy has not been reliable in treatment studies.

CAUTION:

Conditions which can cause a false Negative culture:

1. Patient is on antibiotics. Please repeat culture post therapy.

2. Anaerobic bacterial infection.

3. Fastidious aerobic bacteria which are not able to grow on routine culture media.

4. Besides all these factors, at least in 25-40 % of cases there is no direct correlation between in vivo clinical picture.

5. Renal tuberculosis to be confirmed by AFB studies.

*** End Of Report ***





DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)

NOT VALID FOR MEDICO LEGAL PURPOSE

440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. **REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)**

