

## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. VIKAS GUPTA

**AGE/ GENDER** : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** REG. NO./LAB NO. : 122503200002

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Mar/2025 12:46PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

## **SWASTHYA WELLNESS PANEL: 1.4 COMPLETE BLOOD COUNT (CBC)**

#### **RED BLOOD CELLS (RBCS) COUNT AND INDICES**

HAEMOGLOBIN (HB) by CALORIMETRIC	15.2	gm/dL	12.0 - 17.0
RED BLOOD CELL (RBC) COUNT by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	5.56 <sup>H</sup>	Millions/cmm	3.50 - 5.00
PACKED CELL VOLUME (PCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	44.6	%	40.0 - 54.0
MEAN CORPUSCULAR VOLUME (MCV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	80.1	fL	80.0 - 100.0
MEAN CORPUSCULAR HAEMOGLOBIN (MCH) by Calculated by automated hematology analyzer	27.3	pg	27.0 - 34.0
MEAN CORPUSCULAR HEMOGLOBIN CONC. (MCHC) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	34.1	g/dL	32.0 - 36.0
RED CELL DISTRIBUTION WIDTH (RDW-CV) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	13	%	11.00 - 16.00
RED CELL DISTRIBUTION WIDTH (RDW-SD) by CALCULATED BY AUTOMATED HEMATOLOGY ANALYZER	40.4	fL	35.0 - 56.0
MENTZERS INDEX by CALCULATED	14.41	RATIO	BETA THALASSEMIA TRAIT: < 13.0 IRON DEFICIENCY ANEMIA: >13.0
GREEN & KING INDEX by CALCULATED	18.7	RATIO	BETA THALASSEMIA TRAIT:<= 65.0 IRON DEFICIENCY ANEMIA: > 65.0
WHITE BLOOD CELLS (WBCS)			
TOTAL LEUCOCYTE COUNT (TLC) by flow cytometry by Sf cube & microscopy	9120	/cmm	4000 - 11000
<u>DIFFERENTIAL LEUCOCYTE COUNT (DLC)</u>			
NEUTROPHILS by Flow cytometry by Sf cube & microscopy	54	%	50 - 70
LYMPHOCYTES	33	%	20 - 40



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





CLIENT CODE.



# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

REPORTING DATE

: 20/Mar/2025 12:46PM

NAME : Mr. VIKAS GUPTA

**AGE/ GENDER** : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name	Value	Unit	Biological Reference interval
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
EOSINOPHILS	4	%	1 - 6
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
MONOCYTES	9	%	2 - 12
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY BASOPHILS	0	%	0 - 1
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	U	70	0 - 1
ABSOLUTE LEUKOCYTES (WBC) COUNT			
ABSOLUTE NEUTROPHIL COUNT	4925	/cmm	2000 - 7500
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
ABSOLUTE LYMPHOCYTE COUNT	3010 <sup>L</sup>	/cmm	800 - 4900
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	A. PKR		
ABSOLUTE EOSINOPHIL COUNT by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	365	/cmm	40 - 440
ABSOLUTE MONOCYTE COUNT	821	/cmm	80 - 880
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY	021	7 CHIIII	00 - 000
ABSOLUTE BASOPHIL COUNT	0	/cmm	0 - 110
by FLOW CYTOMETRY BY SF CUBE & MICROSCOPY			
PLATELETS AND OTHER PLATELET PREDICTIVE	MARKERS.		
PLATELET COUNT (PLT)	225000	/cmm	150000 - 450000
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELETCRIT (PCT)	0.27	%	0.10 - 0.36
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE MEAN PLATELET VOLUME (MPV)	U	fL	6.50 - 12.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	12 <sup>H</sup>	IL	0.30 - 12.0
PLATELET LARGE CELL COUNT (P-LCC)	95000 <sup>H</sup>	/cmm	30000 - 90000
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE	33000		
PLATELET LARGE CELL RATIO (P-LCR)	42.2	%	11.0 - 45.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE			
PLATELET DISTRIBUTION WIDTH (PDW)	17.1 <sup>H</sup>	%	15.0 - 17.0
by HYDRO DYNAMIC FOCUSING, ELECTRICAL IMPEDENCE NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD			
NOTE, TEST CONDUCTED ON EDTA WHOLE BLOOD			



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)



## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

**NAME** : Mr. VIKAS GUPTA

AGE/ GENDER : 46 YRS/MALE **PATIENT ID** : 1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Mar/2025 03:33PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

#### **GLYCOSYLATED HAEMOGLOBIN (HBA1C)**

% GLYCOSYLATED HAEMOGLOBIN (HbA1c): 8.9<sup>H</sup> 4.0 - 6.4

WHOLE BLOOD

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

ESTIMATED AVERAGE PLASMA GLUCOSE 208.73<sup>H</sup> 60.00 - 140.00 mg/dL

by HPLC (HIGH PERFORMANCE LIQUID CHROMATOGRAPHY)

#### INTERPRETATION:

AS PER AMERICAN DIABETES ASSOCIATION (ADA):			
REFERENCE GROUP	GLYCOSYLATED HEMOGLO	OGIB (HBAIC) in %	
Non diabetic Adults >= 18 years	<5.7		
At Risk (Prediabetes)	5.7 – 6.4		
Diagnosing Diabetes	>= 6.5		
	Age > 19 Years		
	Goals of Therapy:	< 7.0	
Therapeutic goals for glycemic control	Actions Suggested:	>8.0	
	Age < 19 Years		
	Goal of therapy:	<7.5	

#### COMMENTS:

- 1. Glycosylated hemoglobin (HbA1c) test is three monthly monitoring done to assess compliace with therapeutic regimen in diabetic patients. 2. Since Hb1c reflects long term fluctuations in blood glucose concentration, a diabetic patient who has recently under good control may still have high concentration of HbAlc. Converse is true for a diabetic previously under good control but now poorly controlled.
- 3. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targetting a goal of < 7.0% may not be
- 4.High HbA1c (>9.0 -9.5 %) is strongly associated with risk of development and rapid progression of microvascular and nerve complications 5. Any condition that shorten RBC life span like acute blood loss, hemolytic anemia falsely lower HbA1c results.
- 6.HbA1c results from patients with HbSS,HbSC and HbD must be interpreted with caution, given the pathological processes including anemia, increased red cell turnover, and transfusion requirement that adversely impact HbA1c as a marker of long-term gycemic control.

7. Specimens from patients with polycythemia or post-splenctomy may exhibit increse in HbA1c values due to a somewhat longer life span of the red cells.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





## A PIONEER DIAGNOSTIC CENTRE

REPORTING DATE

**NAME** : Mr. VIKAS GUPTA

AGE/ GENDER : 46 YRS/MALE **PATIENT ID** : 1798784

**COLLECTED BY** REG. NO./LAB NO. : 122503200002

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM : 12507593 BARCODE NO. **COLLECTION DATE** : 20/Mar/2025 08:40AM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Value** Unit **Test Name Biological Reference interval** 

### **ERYTHROCYTE SEDIMENTATION RATE (ESR)**

ERYTHROCYTE SEDIMENTATION RATE (ESR)

mm/1st hr

0 - 20

: 20/Mar/2025 03:56PM

by RED CELL AGGREGATION BY CAPILLARY PHOTOMETRY

#### INTERPRETATION:

CLIENT CODE.

- 1. ESR is a non-specific test because an elevated result often indicates the presence of inflammation associated with infection, cancer and auto-immune disease, but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it.
- 2. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other test such as C-reactive protein
- 3. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as some others, such as systemic lupus erythematosus

#### CONDITION WITH LOW ESR

A low ESR can be seen with conditions that inhibit the normal sedimentation of red blood cells, such as a high red blood cell count (polycythaemia), significantly high white blood cell count (leucocytosis), and some protein abnormalities. Some changes in red cell shape (such as sickle cells in sickle cell anaemia) also lower the ESR.

#### NOTE:

- 1. ESR and C reactive protein (C-RP) are both markers of inflammation.
- 2. Generally, ESR does not change as rapidly as does CRP, either at the start of inflammation or as it resolves.
   3. CRP is not affected by as many other factors as is ESR, making it a better marker of inflammation.
   4. If the ESR is elevated, it is typically a result of two types of proteins, globulins or fibringen.
   5. Women tend to average mathyldone and entraceptives professional processing mathyldone and with the opposition of the oppositio

- 6. Drugs such as dextran, methyldopa, oral contraceptives, penicillamine procainamide, theophylline, and vitamin A can increase ESR, while aspirin, cortisone, and quinine may decrease it



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

**NAME** : Mr. VIKAS GUPTA

AGE/ GENDER : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Mar/2025 12:46PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

## CLINICAL CHEMISTRY/BIOCHEMISTRY **GLUCOSE FASTING (F)**

GLUCOSE FASTING (F): PLASMA NORMAL: < 100.0  $136.37^{H}$ mg/dL

by GLUCOSE OXIDASE - PEROXIDASE (GOD-POD) PREDIABETIC: 100.0 - 125.0

DIABETIC: > 0R = 126.0

INTERPRETATION
IN ACCORDANCE WITH AMERICAN DIABETES ASSOCIATION GUIDELINES:

1. A fasting plasma glucose level below 100 mg/dl is considered normal.

2. A fasting plasma glucose level between 100 - 125 mg/dl is considered as glucose intolerant or prediabetic. A fasting and post-prandial blood

test (after consumption of 75 gms of glucose) is recommended for all such patients.

3. A fasting plasma glucose level of above 125 mg/dl is highly suggestive of diabetic state. A repeat post-prandial is strongly recommended for all such patients. A fasting plasma glucose level in excess of 125 mg/dl on both occasions is confirmatory for diabetic state.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



CLIENT CODE.



# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

REPORTING DATE

: 20/Mar/2025 12:46PM

**NAME** : Mr. VIKAS GUPTA

**AGE/ GENDER** : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** REG. NO./LAB NO. : 122503200002

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name	Value	Unit	Biological Reference interval
	LIPID PROFILE	: BASIC	
CHOLESTEROL TOTAL: SERUM by CHOLESTEROL OXIDASE PAP	240.26 <sup>H</sup>	mg/dL	OPTIMAL: < 200.0 BORDERLINE HIGH: 200.0 - 239.0 HIGH CHOLESTEROL: > OR = 240.0
TRIGLYCERIDES: SERUM by GLYCEROL PHOSPHATE OXIDASE (ENZYMATIC)	154.47 <sup>H</sup>	mg/dL	OPTIMAL: < 150.0 BORDERLINE HIGH: 150.0 - 199.0 HIGH: 200.0 - 499.0 VERY HIGH: > OR = 500.0
HDL CHOLESTEROL (DIRECT): SERUM by SELECTIVE INHIBITION	39.27	mg/dL	LOW HDL: < 30.0 BORDERLINE HIGH HDL: 30.0 - 60.0 HIGH HDL: > OR = 60.0
LDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	170.1 <sup>H</sup>	mg/dL	OPTIMAL: < 100.0 ABOVE OPTIMAL: 100.0 - 129.0 BORDERLINE HIGH: 130.0 - 159.0 HIGH: 160.0 - 189.0 VERY HIGH: > OR = 190.0
NON HDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	200.99 <sup>H</sup>	mg/dL	OPTIMAL: < 130.0 ABOVE OPTIMAL: 130.0 - 159.0 BORDERLINE HIGH: 160.0 - 189.0 HIGH: 190.0 - 219.0 VERY HIGH: > OR = 220.0
VLDL CHOLESTEROL: SERUM by CALCULATED, SPECTROPHOTOMETRY	30.89	mg/dL	0.00 - 45.00
TOTAL LIPIDS: SERUM by CALCULATED, SPECTROPHOTOMETRY	634.99	mg/dL	350.00 - 700.00
CHOLESTEROL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	6.12 <sup>H</sup>	RATIO	LOW RISK: 3.30 - 4.40 AVERAGE RISK: 4.50 - 7.0 MODERATE RISK: 7.10 - 11.0



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)



HIGH RISK: > 11.0





## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

: 20/Mar/2025 12:46PM

NAME : Mr. VIKAS GUPTA

AGE/ GENDER : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. **COLLECTION DATE** : 20/Mar/2025 08:40AM : 12507593

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name	Value	Unit	Biological Reference interval
LDL/HDL RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	4.33 <sup>H</sup>	RATIO	LOW RISK: 0.50 - 3.0 MODERATE RISK: 3.10 - 6.0 HIGH RISK: > 6.0
TRIGLYCERIDES/HDL RATIO: SERUM by CALCULATED. SPECTROPHOTOMETRY	3.93	RATIO	3.00 - 5.00

#### **INTERPRETATION:**

CLIENT CODE.

1. Measurements in the same patient can show physiological analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.

2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.

3. Low HDL levels are associated with increased risk for Atherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available

to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.

4. NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants) along with LDL-cholesterol as co-primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non

5. Additional testing for Apolipoprotein B, hsCRP,Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. VIKAS GUPTA

**AGE/ GENDER** : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Mar/2025 12:46PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Test Name Value** Unit **Biological Reference interval** 

#### **LIVER FUNCTION TEST (COMPLETE)**

BILIRUBIN TOTAL: SERUM by DIAZOTIZATION, SPECTROPHOTOMETRY	0.76	mg/dL	INFANT: 0.20 - 8.00 ADULT: 0.00 - 1.20
BILIRUBIN DIRECT (CONJUGATED): SERUM by DIAZO MODIFIED, SPECTROPHOTOMETRY	0.25	mg/dL	0.00 - 0.40
BILIRUBIN INDIRECT (UNCONJUGATED): SERUM by CALCULATED, SPECTROPHOTOMETRY	0.51	mg/dL	0.10 - 1.00
SGOT/AST: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	31.55	U/L	7.00 - 45.00
SGPT/ALT: SERUM by IFCC, WITHOUT PYRIDOXAL PHOSPHATE	45.27	U/L	0.00 - 49.00
AST/ALT RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	0.7	RATIO	0.00 - 46.00
ALKALINE PHOSPHATASE: SERUM by Para nitrophenyl phosphatase by amino methyl propanol	101	U/L	40.0 - 130.0
GAMMA GLUTAMYL TRANSFERASE (GGT): SERUM by SZASZ, SPECTROPHTOMETRY	27.86	U/L	0.00 - 55.0
TOTAL PROTEINS: SERUM by BIURET, SPECTROPHOTOMETRY	6.54	gm/dL	6.20 - 8.00
ALBUMIN: SERUM by BROMOCRESOL GREEN	4.28	gm/dL	3.50 - 5.50
GLOBULIN: SERUM by CALCULATED, SPECTROPHOTOMETRY	2.26 <sup>L</sup>	gm/dL	2.30 - 3.50
A : G RATIO: SERUM	1.89	RATIO	1.00 - 2.00

#### INTERPRETATION

by CALCULATED, SPECTROPHOTOMETRY

NOTE:- To be correlated in individuals having SGOT and SGPT values higher than Normal Referance Range.

**USE**:- Differential diagnosis of diseases of hepatobiliary system and pancreas.

#### INCREASED:

DRUG HEPATOTOXICITY	> 2
ALCOHOLIC HEPATITIS	> 2 (Highly Suggestive)
CIRRHOSIS	1.4 - 2.0
INTRAHEPATIC CHOLESTATIS	> 1.5
HEPATOCELLULAR CARCINOMA & CHRONIC HEPATITIS	> 1.3 (Slightly Increased)



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST







## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. VIKAS GUPTA

**AGE/ GENDER** : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Mar/2025 12:46PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Test Name Value** Unit **Biological Reference interval** 

#### **DECREASED:**

1. Acute Hepatitis due to virus, drugs, toxins (with AST increased 3 to 10 times upper limit of normal)

2. Extra Hepatic cholestatis: 0.8 (normal or slightly decreased).

#### PROGNOSTIC SIGNIFICANCE:

NORMAL	< 0.65
GOOD PROGNOSTIC SIGN	0.3 - 0.6
POOR PROGNOSTIC SIGN	1.2 - 1.6



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





## PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

### A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

**NAME** : Mr. VIKAS GUPTA

**AGE/ GENDER** : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Mar/2025 04:28PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name	Value	Unit	Biological Reference interval
KIDN	EY FUNCTION T	EST (COMPLETE)	
UREA: SERUM by urease - glutamate dehydrogenase (gldh)	20.54	mg/dL	10.00 - 50.00
CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY	0.95	mg/dL	0.40 - 1.40
BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETRY	9.6	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	10.11	RATIO	10.0 - 20.0

UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETRY	21.62	RATIO	
URIC ACID: SERUM	3.67	mg/dL	3.60 - 7.70

CALCIUM: SERUM 9.49 mg/dL 8.50 - 10.60by ARSENAZO III, SPECTROPHOTOMETRY PHOSPHOROUS: SERUM 2.30 - 4.70 2.49 mg/dL

by PHOSPHOMOLYBDATE, SPECTROPHOTOMETRY

**ELECTROLYTES** SODIUM: SERUM 138.4 mmol/L 135.0 - 150.0 by ISE (ION SELECTIVE ELECTRODE) POTASSIUM: SERUM 4.78 mmol/L 3.50 - 5.00by ISE (ION SELECTIVE ELECTRODE) 103.8 90.0 - 110.0 CHLORIDE: SERUM mmol/L by ISE (ION SELECTIVE ELECTRODE)

#### **ESTIMATED GLOMERULAR FILTERATION RATE**

ESTIMATED GLOMERULAR FILTERATION RATE 100

(eGFR): SERUM by CALCULATED **INTERPRETATION:** 

To differentiate between pre- and post renal azotemia.

#### INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

- 1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.
- Catabolic states with increased tissue breakdown.
- 3. GI haemorrhage.



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

: 20/Mar/2025 04:28PM

**NAME** : Mr. VIKAS GUPTA

AGE/ GENDER : 46 YRS/MALE **PATIENT ID** : 1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Test Name Value** Unit **Biological Reference interval** 

REPORTING DATE

4. High protein intake.

CLIENT CODE.

5. Impaired renal function plus

6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushing's syndrome, high protein diet, burns, surgery, cachexia, high fever).

7. Urine reabsorption (e.g. ureter colostomy)

8. Reduced muscle mass (subnormal creatinine production)

9. Certain drugs (e.g. tetracycline, glucocorticoids)

#### INCREASED RATIO (>20:1) WITH ELEVATED CREATININE LEVELS:

- 1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).
- 2. Prerenal azotemia superimposed on renal disease.

#### DECREASED RATIO (<10:1) WITH DECREASED BUN:

- 1. Acute tubular necrosis.
- 2. Low protein diet and starvation.
- 3. Severe liver disease.
- 4. Other causes of decreased urea synthesis.
- 5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).
- 6. Inherited hyperammonemias (urea is virtually absent in blood).
- 7. SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.
- 8. Pregnancy.

### **DECREASED RATIO (<10:1) WITH INCREASED CREATININE:**

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure.

#### **INAPPROPIATE RATIO:**

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

2. Cephalosporin therapy (interferes with creatinine measurement). **ESTIMATED GLOMERULAR FILTERATION RATE**:

CKD STAGE	DESCRIPTION	GFR ( mL/min/1.73m2 )	ASSOCIATED FINDINGS
G1	Normal kidney function	>90	No proteinuria
G2	Kidney damage with normal or high GFR	>90	Presence of Protein , Albumin or cast in urine
G3a	Mild decrease in GFR	60 -89	
G3b	Moderate decrease in GFR	30-59	
G4	Severe decrease in GFR	15-29	
G5	Kidney failure	<15	



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)







## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

: 20/Mar/2025 04:28PM

**NAME** : Mr. VIKAS GUPTA

AGE/ GENDER : 46 YRS/MALE **PATIENT ID** : 1798784

**COLLECTED BY** REG. NO./LAB NO. : 122503200002

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. **COLLECTION DATE** : 20/Mar/2025 08:40AM : 12507593

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

**Test Name Value** Unit **Biological Reference interval** 

REPORTING DATE

COMMENTS:

CLIENT CODE.

1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a measure of functioning nephrons of the kidney.

2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD

4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure
6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY)





## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

REPORTING DATE

: 20/Mar/2025 04:28PM

**NAME** : Mr. VIKAS GUPTA

**AGE/ GENDER** : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name	Value	Unit	Biological Reference interval
	IRON PROFILE		
IRON: SERUM by FERROZINE, SPECTROPHOTOMETRY	134.8	μg/dL	59.0 - 158.0
UNSATURATED IRON BINDING CAPACITY (UIBC) :SERUM by FERROZINE, SPECTROPHOTOMETERY	205.45	μg/dL	150.0 - 336.0
TOTAL IRON BINDING CAPACITY (TIBC) :SERUM by SPECTROPHOTOMETERY	340.25	μg/dL	230 - 430
%TRANSFERRIN SATURATION: SERUM by CALCULATED, SPECTROPHOTOMETERY (FERENE)	39.62	%	15.0 - 50.0
TRANSFERRIN: SERUM by SPECTROPHOTOMETERY (FERENE)	241.58	mg/dL	200.0 - 350.0

#### INTERPRETATION:

CLIENT CODE.

INTERIALION.			
VARIABLES	ANEMIA OF CHRONIC DISEASE	IRON DEFICIENCY ANEMIA	THALASSEMIA α/β TRAIT
SERUM IRON:	Normal to Reduced	Reduced	Normal
TOTAL IRON BINDING CAPACITY:	Decreased	Increased	Normal
% TRANSFERRIN SATURATION:	Decreased	Decreased < 12-15 %	Normal
SERUM FERRITIN:	Normal to Increased	Decreased	Normal or Increased

#### **IRON**:

1. Serum iron studies is recommended for differential diagnosis of microcytic hypochromic anemia.i.e iron deficiency anemia, zinc deficiency anemia, anemia of chronic disease and thalassemia syndromes.

2. It is essential to isolate iron deficiency anemia from Beta thalassemia syndromes because during iron replacement which is therapeutic for iron deficiency anemia, is severely contra-indicated in Thalassemia.

TOTAL IRON BINDING CAPACITY (TIBC):

1.It is a direct measure of protein transferrin which transports iron from the gut to storage sites in the bone marrow.

% TRANSFERRIN SATURATION:

1.Occurs in idiopathic hemochromatosis and transfusional hemosiderosis where no unsaturated iron binding capacity is available for iron mobilization. Similar condition is seen in congenital deficiency of transferrin.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







## A PIONEER DIAGNOSTIC CENTRE

**NAME** : Mr. VIKAS GUPTA

AGE/ GENDER : 46 YRS/MALE **PATIENT ID** : 1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Mar/2025 04:37PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

### **ENDOCRINOLOGY**

#### THYROID FUNCTION TEST: TOTAL

TRIIODOTHYRONINE (T3): SERUM 0.35 - 1.931.27 ng/mL by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY) THYROXINE (T4): SERUM 8.85 μgm/dL 4.87 - 12.60by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

THYROID STIMULATING HORMONE (TSH): SERUM 2.31 0.35 - 5.50μIU/mL

by CMIA (CHEMILUMINESCENT MICROPARTICLE IMMUNOASSAY)

#### 3rd GENERATION, ULTRASENSITIVE INTERPRETATION:

TSH levels are subject to circadian variation, reaching peak levels between 2-4 a.m and at a minimum between 6-10 pm. The variation is of the order of 50%. Hence time of the day has influence on the measured serum TSH concentrations. TSH stimulates the production and secretion of the metabolically active hormones, thyroxine (T4) and triliodothyronine (T3). Failure at any level of regulation of the hypothalamic-pituitary-thyroid axis will result in either underproduction (hypothyroidism) or overproduction(hyperthyroidism) of T4 and/or T3.

CLINICAL CONDITION	T3	T4	TSH
Primary Hypothyroidism:	Reduced	Reduced	Increased (Significantly)
Subclinical Hypothyroidism:	Normal or Low Normal	Normal or Low Normal	High
Primary Hyperthyroidism:	Increased	Increased	Reduced (at times undetectable)
Subclinical Hyperthyroidism:	Normal or High Normal	Normal or High Normal	Reduced

- 1. T3 and T4 circulates in reversibly bound form with Thyroid binding globulins (TBG), and to a lesser extent albumin and Thyroid binding Pre Albumin so conditions in which TBG and protein levels alter such as pregnancy, excess estrogens, androgens, anabolic steroids and glucocorticoids may falsely affect the T3 and T4 levels and may cause false thyroid values for thyroid function tests
- 2. Normal levels of T4 can also be seen in Hyperthyroid patients with :T3 Thyrotoxicosis, Decreased binding capacity due to hypoproteinemia or ingestion of certain drugs
- 3. Serum T4 levels in neonates and infants are higher than values in the normal adult, due to the increased concentration of TBG in neonate serum.
- 4. TSH may be normal in central hypothyroidism, recent rapid correction of hyperthyroidism or hypothyroidism, pregnancy, phenytoin therapy.

TRIIODOTHY	TRIIODOTHYRONINE (T3)		INE (T4)	THYROID STIMULATING HORMONE (TSH)	
Age	Refferance Range (ng/mL)	Age	Refferance Range (µg/dL)	Age	Reference Range ( μΙU/mL)
0 - 7 Days	0.20 - 2.65	0 - 7 Days	5.90 - 18.58	0 - 7 Days	2.43 - 24.3
7 Days - 3 Months	0.36 - 2.59	7 Days - 3 Months	6.39 - 17.66	7 Days - 3 Months	0.58 - 11.00
3 - 6 Months	0.51 - 2.52	3 - 6 Months	6.75 – 17.04	3 Days – 6 Months	0.70 - 8.40
6 - 12 Months	0.74 - 2.40	6 - 12 Months	7.10 – 16.16	6 – 12 Months	0.70 - 7.00



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. VIKAS GUPTA

AGE/ GENDER : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** REG. NO./LAB NO. : 122503200002

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Mar/2025 04:37PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name			Value	Unit	•	Biolog	ical Reference interval
1 - 10 Years	0.92 - 2.28	1 - 10 Years	6.00 - 13.80	1 – 10 Years	0.60 - 5.50		
11- 19 Years	0.35 - 1.93	11 - 19 Years	4.87- 13.20	11 – 19 Years	0.50 - 5.50		
> 20 years (Adults)	0.35 - 1.93	> 20 Years (Adults)	4.87 - 12.60	> 20 Years (Adults)	0.35-5.50		
	RECOM	MENDATIONS OF TSH LI	VELS DURING PREGI	NANCY ( µIU/mL)			
	1st Trimester			0.10 - 2.50			
	2nd Trimester		0.20 - 3.00				
	3rd Trimester			0.30 - 4.10			

#### **INCREASED TSH LEVELS:**

- 1. Primary or untreated hypothyroidism may vary from 3 times to more than 100 times normal depending upon degree of hypofunction.
- 2. Hypothyroid patients receiving insufficient thyroid replacement therapy.
- 3. Hashimotos thyroiditis
- 4.DRUGS: Amphetamines, iodine containing agents & dopamine antagonist.
- 5. Neonatal period, increase in 1st 2-3 days of life due to post-natal surge

#### **DECREASED TSH LEVELS:**

- 1. Toxic multi-nodular goiter & Thyroiditis.
- 2. Over replacement of thyroid hormone in treatment of hypothyroidism.
- 3. Autonomously functioning Thyroid adenoma
- 4. Secondary pituitary or hypothalamic hypothyroidism
- 5. Acute psychiatric illness
- 6. Severe dehydration.
- 7.DRUGS: Glucocorticoids, Dopamine, Levodopa, T4 replacement therapy, Anti-thyroid drugs for thyrotoxicosis.

8. Pregnancy: 1st and 2nd Trimester



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. VIKAS GUPTA

**AGE/ GENDER** : 46 YRS/MALE **PATIENT ID** : 1798784

**COLLECTED BY** REG. NO./LAB NO. : 122503200002

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM

CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 20/Mar/2025 12:46PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

**Value** Unit **Biological Reference interval Test Name** 

### **CLINICAL PATHOLOGY URINE ROUTINE & MICROSCOPIC EXAMINATION**

#### **PHYSICAL EXAMINATION**

QUANTITY RECIEVED by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	25	ml	
COLOUR by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	PALE YELLOW		PALE YELLOW
TRANSPARANCY by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	CLEAR		CLEAR
SPECIFIC GRAVITY	1.02		1.002 - 1.030

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY			
CHEMICAL EXAMINATION			
REACTION by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	ACIDIC		
PROTEIN by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)
SUGAR by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)
pH by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	5.5		5.0 - 7.5
BILIRUBIN by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)
NITRITE by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY.	NEGATIVE (-ve)		NEGATIVE (-ve)
UROBILINOGEN by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NOT DETECTED	EU/dL	0.2 - 1.0
KETONE BODIES by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)
BLOOD  by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)
ASCORBIC ACID  by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY	NEGATIVE (-ve)		NEGATIVE (-ve)



MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

/HPF

0 - 3



NEGATIVE (-ve)

**MICROSCOPIC EXAMINATION** RED BLOOD CELLS (RBCs)



CLIENT CODE.



# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

REPORTING DATE

: 20/Mar/2025 12:46PM

NAME : Mr. VIKAS GUPTA

**AGE/ GENDER** : 46 YRS/MALE **PATIENT ID** :1798784

**COLLECTED BY** : 122503200002 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 20/Mar/2025 08:23 AM BARCODE NO. : 12507593 **COLLECTION DATE** : 20/Mar/2025 08:40AM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name	Value	Unit	Biological Reference interval
by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT			
PUS CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	3-4	/HPF	0 - 5
EPITHELIAL CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	2-3	/HPF	ABSENT
CRYSTALS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
CASTS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
BACTERIA by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
OTHERS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
TRICHOMONAS VAGINALIS (PROTOZOA) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	ABSENT		ABSENT

End Of Report



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST

