



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

NAME : Mrs. LOVELY SETH

AGE/ GENDER : 43 YRS/FEMALE **PATIENT ID** : 1612686

COLLECTED BY : 122503260006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 26/Mar/2025 09:31 AM BARCODE NO. : 12507716 **COLLECTION DATE** : 26/Mar/2025 11:23AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Mar/2025 01:33PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit Test Name **Biological Reference interval**

HAEMATOLOGY HAEMOGLOBIN (HB)

HAEMOGLOBIN (HB) gm/dL 12.0 - 16.0 9.9L

by CALORIMETRIC

INTERPRETATION:-

Hemoglobin is the protein molecule in red blood cells that carries oxygen from the lungs to the bodys tissues and returns carbon dioxide from the tissues back to the lungs.

A low hemoglobin level is referred to as ANEMIA or low red blood count.

ANEMIA (DECRESED HAEMOGLOBIN):

- 1) Loss of blood (traumatic injury, surgery, bleeding, colon cancer or stomach ulcer)
- 2) Nutritional deficiency (iron, vitamin B12, folate)
- 3) Bone marrow problems (replacement of bone marrow by cancer)
- 4) Suppression by red blood cell synthesis by chemotherapy drugs
- 5) Kidney failure
- 6) Abnormal hemoglobin structure (sickle cell anemia or thalassemia). POLYCYTHEMIA (INCREASED HAEMOGLOBIN):

- 1) People in higher altitudes (Physiological)
- 2) Smoking (Secondary Polycythemia)
- 3) Dehydration produces a falsely rise in hemoglobin due to increased haemoconcentration
- 4) Advanced lung disease (for example, emphysema)
- 5) Certain tumors
- 6) A disorder of the bone marrow known as polycythemia rubra vera,
- 7) Abuse of the drug erythropoetin (Epogen) by athletes for blood doping purposes (increasing the amount of oxygen available to the body by chemically raising the production of red blood cells).

NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)





PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

REPORTING DATE

: 26/Mar/2025 01:33PM

NAME : Mrs. LOVELY SETH

AGE/ GENDER : 43 YRS/FEMALE **PATIENT ID** : 1612686

COLLECTED BY : 122503260006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 26/Mar/2025 09:31 AM BARCODE NO. : 12507716 **COLLECTION DATE** : 26/Mar/2025 11:23AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Value Unit Test Name **Biological Reference interval**

CLINICAL CHEMISTRY/BIOCHEMISTRY

CALCIUM

8.74 CALCIUM: SERUM mg/dL 8.50 - 10.60

by ARSENAZO III, SPECTROPHOTOMETRY

INTERPRETATION:-

CLIENT CODE.

- 1. Serum calcium (total) estimation is used for the diagnosis and monitoring of a wide range of disorders including diseases of bone, kidney, parathyroid gland, or gastrointestinal tract.
- 2. Calcium levels may also reflect abnormal vitamin D or protein levels.
- 3. The calcium content of an adult is somewhat over 1 kg (about 2% of the body weight). Of this, 99% is present as calcium hydroxyapatite in bones and <1% is present in the extra-osseous intracellular space or extracellular space (ECS)
- 4. In serum, calcium is bound to a considerable extent to proteins (approximately 40%), 10% is in the form of inorganic complexes, and 50% is present as free or ionized calcium.

NOTE:-Calcium ions affect the contractility of the heart and the skeletal musculature, and are essential for the function of the nervous system. In addition, calcium ions play an important role in blood clotting and bone mineralization.

HYPOCALCEMIA (LOW CALCIUM LEVELS) CAUSES:-

- 1.Due to the absence or impaired function of the parathyroid glands or impaired vitamin-D synthesis.
- 2. Chronic renal failure is also frequently associated with hypocalcemia due to decreased vitamin-D synthesis as well as hyperphosphatemia and skeletal resistance to the action of parathyroid hormone (PTH).
- 3. NOTE:- A characteristic symptom of hypocalcemia is latent or manifest tetany and osteomalacia.

HYPERCALCEMIA (INCREASE CALCIUM LEVELS) CAUSES:-

- 1.Increased mobilization of calcium from the skeletal system or increased intestinal absorption.
- 2. Primary hyperparathyroidism (pHPT)
- 3. Bone metastasis of carcinoma of the breast, prostate, thyroid gland, or lung

NOTE:-Severe hypercalcemia may result in cardiac arrhythmia.



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)

CLIENT CODE.



PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

: 26/Mar/2025 05:50PM

NAME : Mrs. LOVELY SETH

AGE/ GENDER : 43 YRS/FEMALE **PATIENT ID** : 1612686

COLLECTED BY REG. NO./LAB NO. : 122503260006

REFERRED BY **REGISTRATION DATE** : 26/Mar/2025 09:31 AM BARCODE NO. : 12507716 **COLLECTION DATE** : 26/Mar/2025 11:23AM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Unit Value **Biological Reference interval** Test Name

REPORTING DATE

KIDNEY FUNCTION TEST (BASIC)

UREA: SERUM by UREASE - GLUTAMATE DEHYDROGENASE (GLDH)	59.23 ^H	mg/dL	10.00 - 50.00
CREATININE: SERUM by ENZYMATIC, SPECTROPHOTOMETERY	2.81 ^H	mg/dL	0.40 - 1.20
BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETERY	27.68 ^H	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETERY	9.85 ^L	RATIO	10.0 - 20.0
UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETERY	21.08	RATIO	
URIC ACID: SERUM	5.87	mg/dL	2.50 - 6.80



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

A PIONEER DIAGNOSTIC CENTRE

■ 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

NAME : Mrs. LOVELY SETH

AGE/ GENDER : 43 YRS/FEMALE **PATIENT ID** : 1612686

COLLECTED BY : 122503260006 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 26/Mar/2025 09:31 AM BARCODE NO. : 12507716 **COLLECTION DATE** : 26/Mar/2025 11:23AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 26/Mar/2025 05:50PM

CLIENT ADDRESS : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval**

INTERPRETATION:

Normal range for a healthy person on normal diet: 12 - 20

To Differentiate between pre- and postrenal azotemia. INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.

Ž.Catabolic states with increased tissue breakdown.

3.GI hemorrhage.

4. High protein intake.

5. Impaired renal function plus.

6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushings syndrome, high protein diet,

burns, surgery, cachexia, high fever)

7. Urine reabsorption (e.g. ureterocolostomy)
8. Reduced muscle mass (subnormal creatinine production)
9. Certain drugs (e.g. tetracycline, glucocorticoids)
INCREASED RATIO (pia (PLIN rices diegrapartic particular partic

1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).

2. Prerenal azotemia superimposed on renal disease.

DECREASED RATIO (<10:1) WITH DECREASED BUN:

1.Acute tubular necrosis.

2.Low protein diet and starvation.

3. Severe liver disease.

4. Other causes of decreased urea synthesis.

5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).

6.Inherited hyperammonemias (urea is virtually absent in blood)

7.SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.

8. Pregnancy

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure

INAPPROPIATE RATIO

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

2. Cephalosporin therapy (interferes with creatinine measurement).

*** End Of Report ***

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)