



## A PIONEER DIAGNOSTIC CENTRE

**NAME** : Mr. JAGDISH CHAND

AGE/ GENDER : 68 YRS/MALE **PATIENT ID** : 1691521

**COLLECTED BY** REG. NO./LAB NO. : 122503280020

REFERRED BY **REGISTRATION DATE** : 28/Mar/2025 11:15 AM BARCODE NO. : 12507762 **COLLECTION DATE** : 28/Mar/2025 11:32AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 28/Mar/2025 02:31PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit Test Name **Biological Reference interval** 

# **HAEMATOLOGY** HAEMOGLOBIN (HB)

HAEMOGLOBIN (HB) gm/dL 12.0 - 17.0

by CALORIMETRIC

### **INTERPRETATION:-**

Hemoglobin is the protein molecule in red blood cells that carries oxygen from the lungs to the bodys tissues and returns carbon dioxide from the tissues back to the lungs.

A low hemoglobin level is referred to as ANEMIA or low red blood count.

### **ANEMIA (DECRESED HAEMOGLOBIN):**

- 1) Loss of blood (traumatic injury, surgery, bleeding, colon cancer or stomach ulcer)
- 2) Nutritional deficiency (iron, vitamin B12, folate)
- 3) Bone marrow problems (replacement of bone marrow by cancer)
- 4) Suppression by red blood cell synthesis by chemotherapy drugs
- 5) Kidney failure
- 6) Abnormal hemoglobin structure (sickle cell anemia or thalassemia). POLYCYTHEMIA (INCREASED HAEMOGLOBIN):

- 1) People in higher altitudes (Physiological)
- 2) Smoking (Secondary Polycythemia)
- 3) Dehydration produces a falsely rise in hemoglobin due to increased haemoconcentration
- 4) Advanced lung disease (for example, emphysema)
- 5) Certain tumors
- 6) A disorder of the bone marrow known as polycythemia rubra vera,
- 7) Abuse of the drug erythropoetin (Epogen) by athletes for blood doping purposes (increasing the amount of oxygen available to the body by chemically raising the production of red blood cells).

NOTE: TEST CONDUCTED ON EDTA WHOLE BLOOD



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

**NAME** : Mr. JAGDISH CHAND

**AGE/ GENDER** : 68 YRS/MALE **PATIENT ID** : 1691521

**COLLECTED BY** REG. NO./LAB NO. : 122503280020

REFERRED BY **REGISTRATION DATE** : 28/Mar/2025 11:15 AM BARCODE NO. : 12507762 **COLLECTION DATE** : 28/Mar/2025 11:32AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 28/Mar/2025 04:56PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit **Biological Reference interval** Test Name

## **CLINICAL CHEMISTRY/BIOCHEMISTRY**

### KIDNEY FUNCTION TEST (BASIC)

UREA: SERUM	65.03 <sup>H</sup>	mg/dL	10.00 - 50.00
by UREASE - GLUTAMATE DEHYDROGENASE (GLDH) CREATININE: SERUM	2.25 <sup>H</sup>	mg/dL	0.40 - 1.40
by ENZYMATIC, SPECTROPHOTOMETERY BLOOD UREA NITROGEN (BUN): SERUM by CALCULATED, SPECTROPHOTOMETERY	30.39 <sup>H</sup>	mg/dL	7.0 - 25.0
BLOOD UREA NITROGEN (BUN)/CREATININE RATIO: SERUM	13.51	RATIO	10.0 - 20.0
by CALCULATED, SPECTROPHOTOMETERY			
UREA/CREATININE RATIO: SERUM by CALCULATED, SPECTROPHOTOMETERY	28.9	RATIO	
URIC ACID: SERUM	9.7 <sup>H</sup>	mg/dL	3.60 - 7.70



CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)







# A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

**NAME** : Mr. JAGDISH CHAND

AGE/ GENDER : 68 YRS/MALE **PATIENT ID** : 1691521

**COLLECTED BY** : 122503280020 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 28/Mar/2025 11:15 AM BARCODE NO. : 12507762 **COLLECTION DATE** : 28/Mar/2025 11:32AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 28/Mar/2025 04:56PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Test Name Value Unit **Biological Reference interval** 

INTERPRETATION:

Normal range for a healthy person on normal diet: 12 - 20

To Differentiate between pre- and postrenal azotemia. INCREASED RATIO (>20:1) WITH NORMAL CREATININE:

1. Prerenal azotemia (BUN rises without increase in creatinine) e.g. heart failure, salt depletion, dehydration, blood loss) due to decreased glomerular filtration rate.

Ž.Catabolic states with increased tissue breakdown.

3.GI hemorrhage.

4. High protein intake.

5. Impaired renal function plus.

6. Excess protein intake or production or tissue breakdown (e.g. infection, GI bleeding, thyrotoxicosis, Cushings syndrome, high protein diet,

burns, surgery, cachexia, high fever)

7. Urine reabsorption (e.g. ureterocolostomy)
8. Reduced muscle mass (subnormal creatinine production)
9. Certain drugs (e.g. tetracycline, glucocorticoids)
INCREASED RATIO (pia (PLIN rices diegrapartic particular partic

1. Postrenal azotemia (BUN rises disproportionately more than creatinine) (e.g. obstructive uropathy).

2. Prerenal azotemia superimposed on renal disease.

### DECREASED RATIO (<10:1) WITH DECREASED BUN:

1.Acute tubular necrosis.

2.Low protein diet and starvation.

3. Severe liver disease.

4. Other causes of decreased urea synthesis.

5. Repeated dialysis (urea rather than creatinine diffuses out of extracellular fluid).

6.Inherited hyperammonemias (urea is virtually absent in blood)

7.SIADH (syndrome of inappropiate antidiuretic harmone) due to tubular secretion of urea.

8. Pregnancy

DECREASED RATIO (<10:1) WITH INCREASED CREATININE:

- 1. Phenacimide therapy (accelerates conversion of creatine to creatinine).
- 2. Rhabdomyolysis (releases muscle creatinine).
- 3. Muscular patients who develop renal failure

**INAPPROPIATE RATIO:** 

1. Diabetic ketoacidosis (acetoacetate causes false increase in creatinine with certain methodologies, resulting in normal ratio when dehydration should produce an increased BUN/creatinine ratio).

2. Cephalosporin therapy (interferes with creatinine measurement).

DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)



## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

**NAME** : Mr. JAGDISH CHAND

AGE/ GENDER : 68 YRS/MALE **PATIENT ID** : 1691521

**COLLECTED BY** : 122503280020 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 28/Mar/2025 11:15 AM BARCODE NO. **COLLECTION DATE** : 12507762 : 28/Mar/2025 11:32AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 28/Mar/2025 04:57PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit Test Name **Biological Reference interval** 

### GLOMERULAR FILTERATION RATE (GFR) - ESTIMATED

ESTIMATED GLOMERULAR FILTERATION RATE

mL/min/1.73m2

KIDNEY FAILURE: < 15.0

(eGFR): SERUM

by SPECTROPHOTOMETRY-ENZYMATIC, MDRD CALCULATION

**INTERPRETATION:** 

CKD STAGE	DESCRIPTION	GFR ( mL/min/1.73m2 )	ASSOCIATED FINDINGS
G1	Normal kidney function	>90	No proteinuria
G2	Kidney damage with	>90	Presence of Protein,
	normal or high GFR		Albumin or cast in urine
G3a	Mild decrease in GFR	60 -89	
G3b	Moderate decrease in GFR	30-59	
G4	Severe decrease in GFR	15-29	
G5	Kidney failure	<15	

- 1. Estimated Glomerular filtration rate (eGFR) is the sum of filtration rates in all functioning nephrons and so an estimation of the GFR provides a
- measure of functioning nephrons of the kidney.

  2. eGFR calculated using the 2009 CKD-EPI creatinine equation and GFR category reported as per KDIGO guideline 2012

  3. In patients, with eGFR creatinine between 45-59 ml/min/1.73 m2 (G3) and without any marker of Kidney damage, It is recommended to measure eGFR with Cystatin C for confirmation of CKD 4. eGFR category G1 OR G2 does not fullfill the criteria for CKD, in the absence of evidence of Kidney Damage
- 5. In a suspected case of Acute Kidney Injury (AKI), measurement of eGFR should be done after 48-96 hours of any Intervention or procedure 6. eGFR calculated by Serum Creatinine may be less accurate due to certain factors like Race, Muscle Mass, Diet, Certain Drugs. In such cases, eGFR should be calculated using Serum Cystatin C
  7. A decrease in eGFR implies either progressive renal disease, or a reversible process causing decreased nephron function (eg, severe dehydration).

ADVICE:

KDIGO guideline, 2012 recommends Chronic Kidney Disease (CKD) should be classified based on cause, eGFR category and Albuminuria (ACR) category. GFR & ACR category combined together reflect risk of progression and helps Clinician to identify the individual who are progressing at more rapid rate than anticipated



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)





# PKR JAIN HEALTHCARE INSTITUTE

NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 **■** pkrjainhealthcare@gmail.com

5.0 - 7.5

**NAME** : Mr. JAGDISH CHAND

**AGE/ GENDER** : 68 YRS/MALE **PATIENT ID** : 1691521

**COLLECTED BY** : 122503280020 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 28/Mar/2025 11:15 AM BARCODE NO. : 12507762 **COLLECTION DATE** : 28/Mar/2025 11:32AM CLIENT CODE. : P.K.R JAIN HEALTHCARE INSTITUTE REPORTING DATE : 28/Mar/2025 03:59PM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

Value Unit Test Name **Biological Reference interval** 

## CLINICAL PATHOLOGY URINE ROUTINE & MICROSCOPIC EXAMINATION

### PHYSICAL EXAMINATION

QUANTITY RECIEVED ml

PALE YELLOW COLOUR AMBER YELLOW

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY TRANSPARANCY HAZY **CLEAR** 

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

SPECIFIC GRAVITY 1.01 1.002 - 1.030

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

## CHEMICAL EXAMINATION

REACTION **ACIDIC** by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

**PROTEIN** NEGATIVE (-ve) Trace

**SUGAR** Negative NEGATIVE (-ve)

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

<=5.0by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

**BILIRUBIN** NEGATIVE (-ve) Negative

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY NEGATIVE (-ve) NITRITE Negative

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY.

**UROBILINOGEN** EU/dL 0.2 - 1.0Normal

NEGATIVE (-ve) KETONE BODIES Negative

BLOOD Negative NEGATIVE (-ve)

by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

NEGATIVE (-ve) NEGATIVE (-ve) ASCORBIC ACID by DIP STICK/REFLECTANCE SPECTROPHOTOMETRY

MICROSCOPIC EXAMINATION



pН

CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY)



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)



CLIENT CODE.



# PKR JAIN HEALTHCARE INSTITUTE NASIRPUR, Hissar Road, AMBALA CITY- (Haryana)

## A PIONEER DIAGNOSTIC CENTRE

**■** 0171-2532620, 8222896961 ■ pkrjainhealthcare@gmail.com

REPORTING DATE

: 28/Mar/2025 03:59PM

NAME : Mr. JAGDISH CHAND

**AGE/ GENDER** : 68 YRS/MALE **PATIENT ID** : 1691521

**COLLECTED BY** : 122503280020 REG. NO./LAB NO.

REFERRED BY **REGISTRATION DATE** : 28/Mar/2025 11:15 AM BARCODE NO. : 12507762 **COLLECTION DATE** : 28/Mar/2025 11:32AM

**CLIENT ADDRESS** : NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA

: P.K.R JAIN HEALTHCARE INSTITUTE

Test Name	Value	Unit	Biological Reference interval
RED BLOOD CELLS (RBCs) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)	/HPF	0 - 3
PUS CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	5-7	/HPF	0 - 5
EPITHELIAL CELLS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	2-4	/HPF	ABSENT
CRYSTALS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
CASTS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
BACTERIA by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
OTHERS by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	NEGATIVE (-ve)		NEGATIVE (-ve)
TRICHOMONAS VAGINALIS (PROTOZOA) by MICROSCOPY ON CENTRIFUGED URINARY SEDIMENT	ABSENT		ABSENT

\*\*\* End Of Report



DR.VINAY CHOPRA CONSULTANT PATHOLOGIST MBBS, MD (PATHOLOGY & MICROBIOLOGY) MBBS, MD (PATHOLOGY)

DR.YUGAM CHOPRA CONSULTANT PATHOLOGIST



440 Dated 17.5.2012 u/s 80 G OF INCOME TAX ACT. PAN NO. AAAAP1600. REPORT ATTRACTS THE CONDITIONS PRINTED OVERLEAF (P.T.O.)