NAME	: Mrs. RINKI		
AGE/ GENDER	: 24 YRS/FEMALE	**PATIENT ID**	: 1820931
COLLECTED BY	:	**REG. NO./LAB NO.**	**: 122504070019**
REFERRED BY	:	**REGISTRATION DATE**	: 07/Apr/2025 01:51 PM
BARCODE NO.	: 12507943	**COLLECTION DATE**	: 07/Apr/2025 02:02PM
CLIENT CODE.	: P.K.R JAIN HEALTHCARE INSTITUTE	**REPORTING DATE**	: 07/Apr/2025 02:11PM
CLIENT ADDRESS	: NASIRPUR, HISSAR ROAD, AMBALA CITY - HARYANA		

Test Name	Value	Unit	Biological Reference interval

IMMUNOPATHOLOGY/SEROLOGY
URINE PREGNANCY TEST (UPT)

URINE PREGNANCY TEST (UPT)	WEAKLY POSITIVE		NEGATIVE (-)

by IMMUNOCHROMATOGRAPHY

INTERPRETATION:-
Urine Pregnancy test have 99% sensitivity, hence the result must be co-related with clinical findings and ultrasound report.

COMMENTS:
1. In addition to pregnancy elevated hCG levels have been reported with gestation and non-gestational trophoblastic disease.
2. Very early pregnancy contaning low concentration of hormone in urine can give a negative result. In such cases urine should be retested after proper interval.
3. HCG level remain detectable for several weeks after normal delivery after casearean, spontaneous abortion or therapeutic abortion.
4. Even very high levels of hCG give test results as weak positive or negative. Ectopic pregnancy may also give weak positive results.
5. Urine sample with infections and samples with low specific gravity may not give satisfactory results.

Repeat after 5-6 days or
Advise beta HCG for confirmation

*** End Of Report ***

DR.VINAY CHOPRA
CONSULTANT PATHOLOGIST
MBBS, MD (PATHOLOGY & MICROBIOLOGY)

DR.YUGAM CHOPRA
CONSULTANT PATHOLOGIST
MBBS , MD (PATHOLOGY)