

RAJINDER

PID NO: P542100640909

Age: 67.0 Year(s) Sex: Female



Reference: Dr.SELF

Sample Collected At: DR VINAY KUMAR CHOPRA KOS Diagnostic Lab, 6349/I, Nicholson Road, Ambala Cantt, HRY 133001. Sample Processed At: Metropolis Healthcare Ltd E-21, B1 Mohan Co-op Ind

Estate New Delhi-110044

VID: 54213150464616 Registered On: 12/08/2021 05:24 AM Collected On:

> 12/08/2021 5:24AM Reported On:

12/08/2021 08:11 PM

 Investigation
 Observed Value
 Unit
 Biological Reference Interval

 Tacrolimus
 Below 1.5
 ng/mL
 Refer Interpretation

(EDTA Whole Blood, CMIA)

Medical Remarks: Correlate clinically.

Interpretation:

Kidney Transplant Recipient	
1 Month post transplant	12-15 ngmL
2 to 3 months post transplant	10 - 12 ng/mL
4 to 6 months post transplant	05 - 10 ng/mL
>6 months in selected patients levels	<5 ng/mL, may be sufficient if graft function is stable but immunosuppression is highly variable
In Liver Transplant Recipients	05 to 20 ng/mL

- Tacrolimus results can show variations depending on following reasons-
 - Different Route of administration, Time of collection
 - Time after transplant, Type of allograft.
 - Different amount & time of dosage, Drug distribution which inturn depends on hematocrit & plasma proteins concentration.
 - Different drug company
 - Other co-morbid medical conditions, Concomitant use of immunosuppression & other drugs that inhibit (calcium channel blockers, antifungal agents, some antibiotics) or induce (anticonvulsants, rifampin) metabolism for tacrolimus.
 - Different test methodologies can yield different results, it is advisable to monitor the drug levels using the same testing method & laboratory.

Abbreviation:

CMIA: Chemiluminescence Microparticle Immunoassay

-- End of Report --

Dr. Reema Agrawal MD (Pathology)

Page 1 of 1