Basic Inf	ormation									
Name: Weight: Race: LMP Day: Sample in	Asian		Contact: Birthdate: 1991-08- Twins: Single vil Sender:						Year Ison	
	2023-03-11		Sam	ple NO.:A0064	325	Scan Date:	2023-03-0)3		
Lab:	Lab:		Sample	-		GA:	13+1			
BDP:	1	mm	1	CRL 57.10	mm	NT	1.20	mm		
Assay]			• •		• •				
NO.	Item abbr		Result	Unit N		Reference range				
1	free-ß-HO	CG	54.01	ng/ml	0.74					
2	PAPP-A	A	10000.00	mIU/L	0.95					
3	NT		1.20	mm	0.80					
Risk calculate										
Age risk: 1:727						21-3 syndrome risk				
Parameter: Trisomy21 Risk: 1:24646 Cut Off: (< 1:150) Screening ResultNegative					50 	Age		sk above cut ou risk 1: >10		
						18-3 s	syndrome ri	sk		
Parameter: Trisomy18/13					100					
F	Risk: 1:1127116				×		Ris	sk above cut	off	
Cut	Off: (< 1:3	300)			1200 XSI2 200		Yo	ou risk 1:>10	000	
Screening F	Result:Negati	ve			>5000	Age	50			
Parameter:			Cut Off:		·	Screening Result:				

Advice: Diagnostic results with less risk

Note: *The basic information on the basis of Down's risk assessment in this report is provided at the time of your onsite. When you get this report, please first check whether your relevant information is correct. If there is any discrepancy, please contact your doctor in time, so as to feedback us the correct information and documents, then obtain the correct report. *The high risk and borderline risk of trisomy 21 or trisomy 18 requires further interventional prenatal diagnosis (from fetuses such as villus, amniotic fluid, cord blood, etc.); high risk of neural tube defect (NTD), please go to ultrasound prenatal diagnosis qualified hospitals use ultrasound to exclude.

*The risk of NTD is only calculated at 14-22 weeks.

*The screening result with low risk only shows that the chance of this kind of congenital abnormality in your fetus is less, and the possibility of this kind of abnormality or other abnormalities cannot be completely ruled out. Please consult a doctor in time after you get the report, and the doctor will follow your Risks and other conditions (whether you are older than 35 years old, whether you have had more than one child with other deformities, or have other diseases such as tumors) are comprehensively considered to suggest whether you need to take further examination to confirm the diagnosis.

**This report only can be reference and assistant for doctor , cannot directly give conclusion by this **

Doctor: