Basic Inf	ormation							
Name: Weight: Race: LMP Day:	PAYAL 76.40 Kg Asian formation	Contact: Birthdate: 1997-02- Twins: No Sender:		2-21	-	Gender: ed Due age: le method:	Female 26.13 Year CRL Robinson	
	2023-04-07	Sam	ole NO.:100901	07	Scan Date:	2023-04-1	0	
Lab:		Sample I	•		GA:	12+2	· •	
BDP:		•	CRL 58.70	mm	NT	1.30	mm	
Assay]							
NO.	Item abbr	Result	Unit	MOM	Refer	ence range		
1	free-ß-HCG	27.43	ng/ml	0.63				
2	PAPP-A	3180.00	mIU/L	1.00				
3	NT	1.30	mm	0.85				
Risk calculate								
Age risk: 1:1274					21-3 syndrome risk			
I Cut	neter:Trisomy21 Risk: 1:60791 :Off: (< 1:150) ResultNegative			50 	Age		sk above cut off ou risk 1: >10000	
					18-3 5	syndrome ri	sk	
Parameter: Trisomy18/13				100		.,		
Risk: 1:1651471				200 오		Ris	sk above cut off	
Cut	Off: (< 1:300)					Yo	ou risk 1:>10000	
Screening F	Result:Negative			>5000	Age	50		
Parameter:			Cut Off:		Screening Result:			

Advice: Diagnostic results with less risk

Note: *The basic information on the basis of Down's risk assessment in this report is provided at the time of your onsite. When you get this report, please first check whether your relevant information is correct. If there is any discrepancy, please contact your doctor in time, so as to feedback us the correct information and documents, then obtain the correct report. *The high risk and borderline risk of trisomy 21 or trisomy 18 requires further interventional prenatal diagnosis (from fetuses such as villus, amniotic fluid, cord blood, etc.); high risk of neural tube defect (NTD), please go to ultrasound prenatal diagnosis qualified hospitals use ultrasound to exclude.

*The risk of NTD is only calculated at 14-22 weeks.

*The screening result with low risk only shows that the chance of this kind of congenital abnormality in your fetus is less, and the possibility of this kind of abnormality or other abnormalities cannot be completely ruled out. Please consult a doctor in time after you get the report, and the doctor will follow your Risks and other conditions (whether you are older than 35 years old, whether you have had more than one child with other deformities, or have other diseases such as tumors) are comprehensively considered to suggest whether you need to take further examination to confirm the diagnosis.

**This report only can be reference and assistant for doctor , cannot directly give conclusion by this **

Doctor: