Basic Information	
Name: SARABJEET KAUR	Contact: Gender: Female
Weight: 36.00 Kg	Birthdate: 1993-11-08 Age of EDC: 31.04 Year
Race: Asian	Twins: No GA calc method: CRL Robinson
LMP Day:	Sender:
Sample information	
	ample NO.: 23508411 Scan Date: 2024-05-14
Lab: Sa	mple Date: 2024-05-16 GA: 12+6
BPD: mm (CRL length: 61.70 mm NT length: 2.00 mm
Assay	
NO. Item abbr Result	Unit MOM Reference range
1 free-β-HCG 117.00	ng/ml 1.71
2 PAPP-A 10000.00	mIU/L 1.08
3 NT 2.00	mm 1.25
isk calculate	
Age risk: 1:848	21-3 syndrome risk
Parameter: Trisomy21	50
Farameter: msomy21	بن المعنى الم
Risk: 1:2238	You risk 1:2238
Cut Off: (< 1:150)	
Screaning Result: Negative	>5000
Serearing result. Reguive	Age
	18-3 syndrome risk
Parameter: Trisomy18/13	100
Risk: 1:4989761	Risk above cut off
Cut Off: (< 1:300)	$\frac{1}{200}$ 200 - Kisk above cut off You risk 1: >10000
Screening Result: Negative	>5000
	Age
	5

Advice: Diagnostic results with less risk

Note: *The basic information on the basis of Down's risk assessment in this report is provided at the time of your onsite. When you get this report, please first check whether your relevant information is correct. If there is any discrepancy, please contact your doctor in time, so as to feedback us the correct information and documents, then obtain the correct report. *The high risk and borderline risk of trisomy 21 or trisomy 18 requires further interventional prenatal diagnosis (from fetuses such as villus, amniotic fluid, cord blood, etc.); high risk of neural tube defect (NTD), please go to ultrasound prenatal diagnosis qualified hospitals use ultrasound to exclude.

*The risk of NTD is only calculated at 14-22 weeks.

*The screening result with low risk only shows that the chance of this kind of congenital abnormality in your fetus is less, and the possibility of this kind of abnormality or other abnormalities cannot be completely ruled out. Please consult a doctor in time after you get the report, and the doctor will follow your Risks and other conditions (whether you are older than 35 years old, whether you have had more than one child with other deformities, or have other diseases such as tumors) are comprehensively considered to suggest whether you need to take further examination to confirm the diagnosis.

**This report only can be reference and assistant for doctor , cannot directly give conclusion by this **

Doctor: