Basic Infor	mation				
Name: S Weight: 4	SANNJH		Contact: Birthdate: 2005-12-2 Twins: No Sender:	29	Gender: Female Age of EDC: 19.61 Year GA calc method: CRL Robinson
Sample info					
Send time: 2	025-02-01	5	Sample NO.: 10528455	5	Scan Date: 2025-01-31
Lab:		Sa	ample Date: 2025-01-3	1	GA: 12+6
BPD:	mm	(	CRL length: 65.70 n	nm	NT length: 1.70 mm
Assay –					
NO.	Item abbr	Result	Unit	MOM	Reference range
1 fr	ee-β-HCG	98.10	ng/ml	1.66	
2	PAPP-A	7397.00	mIU/L	0.97	
3	NT	1.70	mm	1.01	
isk calculate –					
Age ri	sk: 1:1542				21-3 syndrome risk
Ris Cut O	ter:Trisomy21 sk: 1:8119 off: ( < 1:150 ) sult: Negative			50 ස් 100 >5000	Risk above cut off You risk 1:8119
C	C				
Parameter: Trisomy18/13			100	18-3 syndrome risk	
Risk: 1:12493232			×	Risk above cut off	
Cut Off: ( < 1:300 )			· 200 · 관	You risk 1: >10000	
Screening Res	sult: Negative			>5000	50
					Age

Advice: Diagnostic results with less risk

Note: \*The basic information on the basis of Down's risk assessment in this report is provided at the time of your onsite. When you get this report, please first check whether your relevant information is correct. If there is any discrepancy, please contact your doctor in time, so as to feedback us the correct information and documents, then obtain the correct report. \*The high risk and borderline risk of trisomy 21 or trisomy 18 requires further interventional prenatal diagnosis (from fetuses such as villus, amniotic fluid, cord blood, etc.); high risk of neural tube defect (NTD), please go to ultrasound prenatal diagnosis qualified hospitals use ultrasound to exclude.

\*The risk of NTD is only calculated at 14-22 weeks.

\*The screening result with low risk only shows that the chance of this kind of congenital abnormality in your fetus is less, and the possibility of this kind of abnormality or other abnormalities cannot be completely ruled out. Please consult a doctor in time after you get the report, and the doctor will follow your Risks and other conditions (whether you are older than 35 years old, whether you have had more than one child with other deformities, or have other diseases such as tumors) are comprehensively considered to suggest whether you need to take further examination to confirm the diagnosis.

\*\*This report only can be reference and assistant for doctor , cannot directly give conclusion by this \*\*

Doctor: